

Antidepressant	Advantages During Pregnancy	Teratogenicity	Other Disadvantages During Pregnancy	Estimated % of Maternal Dose to Breastfeeding Baby	Reported Side Effects to Breastfeeding Babies
Bupropion (Wellbutrin: Zyban)	<ul style="list-style-type: none"> Fewer sexual side effects Less risk of weight gain Helps with smoking cessation 	Morphologic - limited evidence of cardiac malformations; increased risk for pulmonary hypertension Behavioral - limited evidence of increased risk of ADHD	<ul style="list-style-type: none"> Limited data available Lowers seizure threshold Can cause insomnia May increase risk of miscarriage 	2.0%	Seizures
Citalopram (Celexa)	<ul style="list-style-type: none"> Few interactions with other medications 	Morphologic - risk of neural tube defect Behavioral - none found	<ul style="list-style-type: none"> Limited data available 	0.7% - 9.0%	Uneasy sleep, drowsiness, irritability, weight loss
Desipramine (Norpramin)	<ul style="list-style-type: none"> More studies in human pregnancy, including neurodevelopmental follow-up 	Morphologic - none found Behavioral - none found	<ul style="list-style-type: none"> Maternal side effects additive to pregnancy effects (sedation, constipation, tachycardia) Orthostatic hypotension, risking decreased placental perfusion Fetal and neonatal side effects: tachycardia, urinary retention 	1.0%	Agitation of newborn, potential triggering of seizure activity if there is a history of seizures
Duloxetine (Cymbalta)	<ul style="list-style-type: none"> Also treats diabetic peripheral neuropathic pain 	Morphologic - unknown Behavioral - unknown	<ul style="list-style-type: none"> No systematic studies in human pregnancy 	0.1%	Unknown
Escitalopram (Lexapro)	<ul style="list-style-type: none"> Few interactions with other medications 	Morphologic - unknown Behavioral - unknown	<ul style="list-style-type: none"> No systematic studies in human pregnancy 	3.9% - 7.9%	Enterocolitis
Fluoxetine (Prozac)	<ul style="list-style-type: none"> More studies in human pregnancy, including metaanalysis and neurodevelopmental follow-up 	Morphologic - increased risk of cardiovascular malformations* Behavioral - none found	<ul style="list-style-type: none"> More reports of neonatal side effects than most other antidepressants 	1.2% - 12.0%	Excessive crying, irritability, vomiting, watery stools, difficulty sleeping, tremor, somnolence, hypotonia, decreased weight gain, hyperglycemia
Mirtazapine (Remeron)	<ul style="list-style-type: none"> Fewer sexual side effects Helps restore appetite in women who are not gaining weight Less likely to exacerbate nausea and vomiting 	Morphologic - none found Behavioral - unknown	<ul style="list-style-type: none"> Limited data available Can cause excessive weight gain Tends to be sedating May increase risk of preterm birth 	0.6% - 2.8%	None
Nortriptyline (Pamelor)	<ul style="list-style-type: none"> More studies in human pregnancy, including neurodevelopmental follow-up 	Morphologic - none found Behavioral - none found	<ul style="list-style-type: none"> Maternal side effects additive to pregnancy effects (sedation, constipation, tachycardia) Orthostatic hypotension, risking decreased placental perfusion Fetal and neonatal side effects: tachycardia, urinary retention 	1.3%	None
Paroxetine (Paxil)	<ul style="list-style-type: none"> Minimal association with cardiovascular malformations but may be optimal for some individual parents 	Morphologic - possible increased risk of cardiovascular malformations Behavioral - unknown	<ul style="list-style-type: none"> More reports of neonatal side effects than most other antidepressants ACOG recommends fetal echo for all exposed fetuses 	0.1% - 4.3%	Irritability, sleepiness, constipation, SIADH
Sertraline (Zoloft)	<ul style="list-style-type: none"> Relatively well-studied in human pregnancy Fewer reports of neonatal side effects than other antidepressants 	Morphologic - unlikely increased risk of omphalocele and septal defects* Behavioral - none found	<ul style="list-style-type: none"> Minimal association with omphalocele and septal defects 	0.4% - 2.3%	Commonly prescribed by OBs and pediatricians
Venlafaxine (Effexor)	<ul style="list-style-type: none"> None specific, but may be optimal for some individual patients 	Morphologic - none found Behavioral - unknown	<ul style="list-style-type: none"> Limited data available 	5.2% - 7.6%	Decreased weight gain
Desvenlafaxine (Pristiq)	<ul style="list-style-type: none"> None specific, but may be optimal for some individual patients 	Morphologic - unknown Behavioral - unknown	<ul style="list-style-type: none"> No systematic studies in human pregnancy 	Unknown	Unknown

* Findings from one study at variance with other data, perhaps due to methodological flaws

Information for Providers on Antidepressants during Pregnancy & Breastfeeding – April 2015

This chart is produced by the University of Illinois at Chicago (UIC) as a summary of research on antidepressants in human pregnancy and breastfeeding.

Sources of data:

- **Pregnancy data:** Data summarized here are from controlled studies in human pregnancy. The Food and Drug Administration (FDA) Pregnancy Risk Categories, as found in the Physicians' Desk Reference¹, are based on both animal and human studies. No antidepressants are yet specifically FDA-approved for use during pregnancy. All antidepressants cross the placenta, so there are none that are 'Category A' ("no risk"). Medications that are non-teratogenic in animal studies but have never been studied in humans are classified as 'Category B'. Since teratogenicity does not generalize across species, a 'Category B' classification does not imply greater safety in human pregnancy than a 'Category C' or 'D' classification. Several medications have been shifted from 'Category B' to 'Category C' or 'Category D' as their risks became better known.
- **Breastfeeding data:** Data about antidepressant effects on breastfeeding babies are predominantly from case reports and case series. For medications with no reported side effects, that does not necessarily mean the medication is "safe"; often it means there are few case reports available. Reported percent of maternal dose to breastfeeding babies are weight-adjusted estimates that include the agent and its active metabolite(s). *Specific references are available on request.

General guideline:

- Optimal treatment is based on individual patient characteristics and clinical judgment, especially weighing medication risks against risks of untreated illness. Risks of untreated perinatal depression may include preterm birth and other obstetric complications, increased risk of infection and difficult temperament in the infant, impaired parenting, and psychological effects such as impaired cognitive development, emotional and behavioral problems and increased reactivity to stress in children.

Antidepressants as a group may be associated with following risks:

- Increased risk of preterm birth and lower gestational age at birth, but without adverse effects on birth weight or Apgar scores • Increased risk of miscarriage, but rates within norms of the general population.

SSRI antidepressants as a group (citalopram, escitalopram, fluoxetine, paroxetine, sertraline) may be associated with the following risks:

- Neonatal side effects, including respiratory distress, excessive crying, changes in sleep and behavioral state, difficulty sleeping, increased or decreased muscle tone, hyperreflexia, seizures, and/or cardiac arrhythmias.
- Most studies have found no increased risk of gestational hypertension. One retrospective study² found a possible increased risk of gestational hypertension.
- Possible increased risk of persistent pulmonary hypertension in the newborn with exposure later in pregnancy.
- Most studies have found no increased risk of birth defects. One retrospective study³ found a possible increased risk of anencephaly, craniosynostosis, and omphalocele; another⁴ found an increased risk of anomalies in general, although absolute risks were small.
- Delay in lactation, however the delay was only for 14 hours on average.
- Kaiser Study showed 2-fold increased risk for Autism spectrum disorder with use of SSRI within one year of delivery and 3-fold increased risk with SSRI use in first trimester.

For questions, call Illinois DocAssist at 1-866-986-2778 or visit www.docassistillinois.org

1. Physician's Desk Reference. Thomson Reuters. Montvale, NJ. 2. Toh et al. Selective serotonin reuptake inhibitor use and risk of gestational hypertension. *Am J Psychiatry*. 2009 Mar;166(3):320-8. 3. Alwan, S. et al. Use of selective serotonin-reuptake inhibitors in pregnancy and the risk of birth defects. *N Engl J Med*. 2007 Jun 28; 356(26):2684-92. 4. Wogelius et al. Maternal use of selective serotonin reuptake inhibitors and risk of congenital malformations. *Epidemiology*. 2006 Nov;17(6):701-4. 5. Suri et al. Effects of Antenatal Depression and Antidepressant treatment on gestational age at birth and risk of preterm birth. *Am J Psychiatry*. 2007 Aug; 164:1206-1213. 6. Figueroa. Use of antidepressants during pregnancy and risk of Attention-Deficit/Hyperactivity Disorder in the offspring. *JDBP*. 2010 Oct. Vol 31, No.8. 7. Alwan et al. Maternal use of Bupropion and risk of congenital heart defects. *Am J Obstet Gynecol* 2010. 8. Kim et al. Pharmacotherapy of postpartum depression: an update. *NIH Expert Opin Pharmacother*. 2014 June; 15(9): 1223-1234.

© 2011 The Board of Trustees of the University of Illinois. All rights reserved. Originally created by the Illinois Perinatal Mental Health Project; revised in 2015 by Illinois DocAssist. **PLEASE NOTE:** This medication chart offers information to healthcare providers regarding the use of antidepressants in pregnant and postpartum women. The Illinois DocAssist consultation service and this chart have been created as general information only and should not be construed as direct advice about managing any particular patient's care. The information in this chart is intended for the provider ONLY and should NOT be shared directly with a patient. Since Illinois DocAssist consultants have no direct knowledge of the patient, they cannot provide the chart information in a manner that is appropriate for an individual patient's needs. **Please use this medication chart for provider information purposes only.**